

To appear in Astrophysical Journal

Radio Emission from SN 2001gd in NGC 5033

Christopher J. Stockdale¹

*Naval Research Laboratory, Code 7213, Washington, DC 20375-5320;
stockdale@nrl.navy.mil*

Kurt W. Weiler

*Naval Research Laboratory, Code 7213, Washington, DC 20375-5320;
Kurt.Weiler@nrl.navy.mil*

Schuyler D. Van Dyk

IPAC/Caltech, Mail Code 100-22, Pasadena, CA 91125; vandyk@ipac.caltech.edu

Marcos J. Montes

*Naval Research Laboratory, Code 7212, Washington, DC 20375-5320;
Marcos.Montes@nrl.navy.mil*

Nino Panagia²

*Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218;
panagia@stsci.edu*

Richard A. Sramek

*National Radio Astronomy Observatory, P.O. Box 0, Socorro, NM 87801;
dsramek@nrao.edu*

M. A. Perez-Torres

*Instituto de Astrofísica de Andalucía, CSIC, Apdo. Correos 3004, 18080 Granada, Spain;
torres@iaa.es*

and

J. M. Marcaide

Depto. de Astronomia Univ. Valencia, 46100 Burjassot, Spain; J.M.Marcaide@uv.es

ABSTRACT

We present the results of monitoring the radio emission from the Type IIb supernova SN 2001gd between 2002 February 8 and 2002 October 28. Most of the data were obtained using the Very Large Array at the five wavelengths of $\lambda\lambda$ 1.3 cm (22.4 GHz), 2.0 cm (14.9 GHz), 3.6 cm (8.44 GHz), 6.2 cm (4.86 GHz), and 21 cm (1.4 GHz). Observations were also made with Giant Meterwave Radio Telescope at λ 21 cm (1.4 GHz). The object was discovered optically well after maximum light, making any determination of the early radio evolution difficult. However, subsequent observations indicate that the radio emission has evolved regularly in both time and frequency and is well described by the SN shock/circumstellar medium interaction model.

Subject headings: galaxies: individual (NGC 5033) – radio continuum: stars – stars: mass-loss – supernovae: general – supernovae: individual (SN 2001gd)

1. Introduction

Supernova (SN) 2001gd in NGC 5033 was discovered at magnitude, $16^m.5$ on 2001 November 24.820 UT by Nakano *et al.* (2001) at RA(J2000) = $13^h13^m23^s.89$, Dec(J2000) = $+36^\circ38'17''.7$, with independent discoveries by other observers following quickly. No source had been detected at that position (limiting mag. ~ 18.5) on a previous image taken on 2001 May 8 and, nothing is visible on the Palomar Sky Survey red and blue plates (Nakano *et al.* 2001). Ten days later, Matheson *et al.* (2001) confirmed the discovery as a type IIb supernova well past maximum light, stating “the spectrum is almost identical to one of SN 1993J obtained on day 93 after explosion.” Using this suggestion to constrain the explosion date for the source, the optical discovery occurred 83 days after the explosion (Nakano *et al.* 2001) and; the first radio observation was 158 days after the explosion. Tully (1988) determined a distance of 18.7 Mpc (assuming $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$) to NGC 5033. Accepting the Tully (1988) distance and using $H_0 = 65 \text{ km s}^{-1} \text{ Mpc}^{-1}$, we will adopt the distance to the SN of 21.6 Mpc, which results in an absolute visible magnitude at discovery of < -15 , although it is not known what the magnitude was at maximum optical brightness (Nakano *et al.* 2001). We use the Tully distance method to maintain uniformity with our distance determinations to other radio SNe including SN 1985L (Van Dyk *et al.* 1998).

¹National Research Council Postdoctoral Fellow.

²On assignment from the Astrophysics Division, Space Science Department of ESA.

Stockdale *et al.* (2002) detected radio emission from SN 2001gd on 2002 February 8.54 with the Very Large Array (VLA)³ at 22.485 GHz (λ 1.3 cm), 8.435 GHz (λ 3.6 cm), 4.885 GHz (λ 6.1 cm), and 1.465 GHz (λ 21 cm). The radio position of RA(J2000) = $13^h 13^m 23^s.90$, Dec(J2000) = $+36^\circ 38' 18''.1$, with an uncertainty of $\pm 0''.2$ in each coordinate (Stockdale *et al.* 2002), is in agreement with the optical position to within the errors. Since this first detection, we have made, and will continue to make, radio observations with the VLA. We note that NGC 5033 is also the host galaxy of two other historical supernovae, SNe 1950C and 1985L (Kowal, Sargent, & Zwicky 1970; Aksenov *et al.* 1985). The latter was also detected in the radio in 1986 (Van Dyk *et al.* 1998). Stockdale *et al.* (2002) did not detect radio emission from either of these historical supernovae, nor do we in any of our subsequent observations.

2. Radio Observations

The VLA radio observations reported here were made between 2002 February 8 and 2002 October 28. VLA phase and flux density calibration and data reduction followed standard procedures (Weiler *et al.* 1986; Weiler, Panagia, & Sramek 1990), using 3C 286 as the primary flux density calibrator and J1310+323 as the phase calibrator and as a secondary flux density calibrator.

In Table 1, we list the flux density observations of the secondary calibrator source J1310+323, which are plotted in Figure 1. The defined position for J1310+323 for phase reference is RA(J2000) = $13^h 10^m 28^s.66$, Dec(J2000) = $+32^\circ 20' 43''.8$.

In Table 2, we list the flux density measurements for SN 2001gd and Figure 2 plots the data. The data appear to be well described by a parameterized model described below and illustrated in Figure 2 .

The flux density measurement error given for the VLA measurements in Table 2 is a combination of the rms map error, which measures the contribution of small unresolved fluctuations in the background emission and random map fluctuations due to receiver noise, and a basic fractional error, ϵ , included to account for the normal inaccuracy of VLA flux density calibration and possible deviations of the primary calibrator from an absolute flux density scale. The final errors (σ_f) as listed in Table 2 are taken as

$$\sigma_f^2 \equiv (\epsilon S_0)^2 + \sigma_0^2 \quad (1)$$

³The VLA telescope of the National Radio Astronomy Observatory is operated by Associated Universities, Inc. under a cooperative agreement with the National Science Foundation.

where S_0 is the measured flux density, σ_0 is the map rms for each observation, and $\epsilon = 0.10$ at 21 cm, 0.05 at 6 cm, 0.05 at 3.6 cm, 0.075 at 2 cm, and 0.10 at 1.2 cm.

3. Discussion

Common properties of radio SNe (RSNe) include non-thermal synchrotron emission with high brightness temperature, turn-on delay at longer wavelengths, power-law decline after maximum with index β , and spectral index α asymptotically decreasing to an optically thin value (Weiler *et al.* 1986). Weiler *et al.* (1986, 1990) have shown that the “mini-shell” model of Chevalier (1982a; 1982b), adequately describes previously known RSNe. In this model, the relativistic electrons and enhanced magnetic fields necessary for synchrotron emission are generated by the SN shock interacting with a relatively high-density envelope of circumstellar material (CSM) which has been ionized and heated by the initial UV/X-ray flash. This dense cocoon is presumed to have been established by a constant mass-loss ($\dot{M}$) rate, in a constant velocity wind (w_{wind} ; i. e., $\rho_{\text{CSM}} \propto \frac{\dot{M}}{w_{\text{wind}} r^2}$) from a red supergiant (RSG) SN progenitor or companion. This ionized CSM is also the source of the initial absorption. The rapid rise in radio flux density results from the shock overtaking progressively more of the wind matter, leaving less of it along the line-of-sight to the observer to absorb the more slowly decreasing synchrotron emission from the shock region.

3.1. Parameterized Model

Following Weiler *et al.* (1986; 1990; 2001; 2002), Montes, Weiler, & Panagia (1997), and Chevalier (1998), we adopt the parameterized model,

$$S(\text{mJy}) = K_1 \left(\frac{\nu}{5 \text{ GHz}} \right)^\alpha \left(\frac{t - t_0}{1 \text{ day}} \right)^\beta e^{-\tau_{\text{external}}} \left(\frac{1 - e^{-\tau_{\text{CSM}_{\text{clumps}}}}}{\tau_{\text{CSM}_{\text{clumps}}}} \right) \left(\frac{1 - e^{-\tau_{\text{internal}}}}{\tau_{\text{internal}}} \right) \quad (2)$$

with

$$\tau_{\text{external}} = \tau_{\text{CSM}_{\text{homogeneous}}} + \tau_{\text{distant}} \quad (3)$$

where

$$\tau_{\text{CSM}_{\text{homogeneous}}} = K_2 \left(\frac{\nu}{5 \text{ GHz}} \right)^{-2.1} \left(\frac{t - t_0}{1 \text{ day}} \right)^\delta, \quad (4)$$

$$\tau_{\text{CSM}_{\text{clumps}}} = K_3 \left(\frac{\nu}{5 \text{ GHz}} \right)^{-2.1} \left(\frac{t - t_0}{1 \text{ day}} \right)^{\delta'}, \quad (5)$$

and

$$\tau_{\text{distant}} = K_4 \left(\frac{\nu}{5 \text{ GHz}} \right)^{-2.1}, \quad (6)$$

with K_1 , K_2 , K_3 , and K_4 corresponding, formally, to the flux density (K_1), homogeneous absorption (K_2 , K_4), and clumpy (or filamentary) absorption (K_3) at 5 GHz one day after the explosion date t_0 . The terms $\tau_{\text{CSM}_{\text{homogeneous}}}$ and $\tau_{\text{CSM}_{\text{clumps}}}$ describe the attenuation of a local homogeneous and clumpy CSM that are near enough to the supernova progenitor that they are affected by the expanding blastwave. The $\tau_{\text{CSM}_{\text{homogeneous}}}$ absorption term is due to an ionized medium which uniformly covers the emitting source. The $\tau_{\text{CSM}_{\text{clumps}}}$ absorption term describes the attenuation produced by an inhomogeneous medium (see Natta & Panagia (1984) for a more detailed discussion). The τ_{distant} term describes the attenuation produced by a homogeneous medium that uniformly covers the source but is so far from the SN progenitor that it is not affected by the blastwave and is constant in time.

All external and clumpy absorbing media are assumed to be purely thermal, singly ionized gas that absorbs via free-free (f-f) transitions with frequency dependence $\nu^{-2.1}$ in the radio. The parameter δ describes the time dependence of the optical depth for the local, uniform medium. For an undecelerated SN shock, $\delta = -3$ is appropriate⁴, and we have assumed this value for our parameterization since the data are insufficient to properly determine δ at this time. The parameter δ' describes the time dependence of the optical depth for the non-uniform (clumpy) absorbing medium. For this source and at this epoch, we neglect the (K_4) term because any distant, time-independent absorption effects will likely be evident only at lower frequencies and later times than are available in our current data set (see e.g. Montes *et al.* 1997; Schlegel *et al.* 1999 for SN 1978K).

Chevalier (1998) proposed that synchrotron self-absorption (SSA) may be a significant source of absorption at high frequencies and early times for some RSNs. Given the relatively poor sampling of the radio emission turn-on at early times, SSA is not expected to be significant and our parameterization of SN 2001gd is not improved by the inclusion of SSA component.

A search for the best parameter fit to Equations 2 through 6 was carried out by minimizing the reduced chi-squared (χ_{red}^2). The best-fit parameter values and comparisons with the type IIb SN 1993J and the other recent RSN in NGC 5033, SN 1985L, are listed in Table 3. The resulting model curves are plotted in Figure 2. The errors listed for the parameter values, determined by the “bootstrap” method, are at the 1σ uncertainty level.

The fitting procedure and examination of Figure 2 indicates that the gross properties of the radio emission from SN 2001gd are relatively well described by a purely thermal absorption model such as that given by Eqs. 2 — 6, with parameter values listed in Table 3.

⁴Weiler *et al.* (1986) show that the absorption term $\tau_{\text{CSM}_{\text{homogeneous}}} \propto r^{-3}$. Since $r \propto t^1$ for an undecelerated shock, $\tau_{\text{CSM}_{\text{homogeneous}}} \propto t^{-3}$ and from Equation 4, it follows that $\delta = -3$ in this case.

The requirement for non-zero K_2 and K_3 parameter values implies the existence of a clumpy or filamentary absorption component embedded in a relatively homogeneous medium (on a more detailed level, this is consistent with Case 2 discussed by Weiler *et al.* 2001, 2002). The distant absorption term (K_4) was set to zero for reasons discussed previously. The parameter values are typical of values determined for other type II SNe, although they are not tightly constrained by the relatively sparse data set.

3.2. Comparison to Other RSNe

It is useful to make distance independent comparisons between SN 2001gd to SN 1985L, also in NGC 5033. Van Dyk *et al.* (1998) reported only two 6 cm detections of SN 1985L approximately 1 year after explosion, with numerous non-detections. SN 1985L was identified as type IIL by Filippenko & Sargent (1985) and Kris (1985). Despite the relatively small amount of radio data available for the SN 1985L, there are some valid comparisons that can be made. SN 2001gd ($L_{6 \text{ cm peak}} = 2.91 \times 10^{27} \text{ erg s}^{-1} \text{ Hz}^{-1}$) is almost ten times more luminous at the 6 cm peak flux density than SN 1985L ($L_{6 \text{ cm peak}} = 3.10 \times 10^{26} \text{ erg s}^{-1} \text{ Hz}^{-1}$; Van Dyk *et al.* 1998).⁵ The 6 cm peak luminosity of SN 2001gd also lies within a range of other type II SNe, for example, SN 1979C ($2.55 \times 10^{27} \text{ erg s}^{-1} \text{ Hz}^{-1}$) and SN 1980K ($1.18 \times 10^{26} \text{ erg s}^{-1} \text{ Hz}^{-1}$). SNe 2001gd and 1985L also have very different times from explosion to 6 cm peak, although in both cases there is great uncertainty in this time scale. SN 2001gd has a much steeper spectral index ($\alpha = -1.4$) than any other well-observed RSN. Weiler *et al.* (2002) demonstrate that most type II SNe have a spectral index ($\alpha > -1.0$), similar to SN 1985L which has a moderate value of $\alpha = -0.58$. The power law decline index for SN 2001gd ($\beta \sim -0.96$) is comparable to that of other type II RSNe.

It is also useful to make some comparisons between SNe 2001gd and 1993J since both are type IIb SNe and SN 1993J was much better sampled than SN 2001gd. Both have relatively equivalent 6 cm peak luminosities, as reported in Table 3. The time to 6 cm peak for SN 1993J is in relatively close agreement with SN 2001gd (~ 170 days), although this value is somewhat atypical when compared with other type II SNe. Type Ib/Ic SNe typically achieve their 6 cm peak luminosity in <40 days, while most type II SNe take from as a few as ~ 100 days to well over 1 year (Weiler *et al.* 2002). Both SNe 2001gd and 1993J have spectral indices < -1.0 , which is more typical of type Ib/Ic SNe (Weiler *et al.* 2002). But,

⁵We have adjusted the luminosity of SN 1985L to account for a Tully (1988) distance to NGC 5033 of 21.6, assuming $H_0 = 65 \text{ km s}^{-1} \text{ Mpc}^{-1}$, Van Dyk *et al.* (1998) assumed $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ for a Tully (1988) distance of 20 Mpc.

SN 2001gd appears to have a power-law decline index comparable to that of SN 1993J and other type II SNe, unlike values reported for type Ib SNe 1983N and 1984L ($\beta \sim -1.6$; Sramek, Panagia, & Weiler 1984; Weiler *et al.* 1986; Panagia, Sramek, & Weiler 1986).

3.3. Estimates of Mass-loss Rates

Using the derived parameter values from the radio light curves, we can estimate the presupernova mass loss rate of the SN 2001gd progenitor star (Weiler, Panagia, & Montes, 2001; Weiler *et al.* 1986; 2001; 2002), as

$$\begin{aligned} \frac{\dot{M}/(M_{\odot} \text{ yr}^{-1})}{w_{\text{wind}}/(10 \text{ km s}^{-1})} &= 3.0 \times 10^{-6} \langle \tau_{\text{eff}}^{0.5} \rangle m^{-1.5} \left(\frac{v_i}{10^4 \text{ km s}^{-1}} \right)^{1.5} \\ &\times \left(\frac{t_i}{45 \text{ days}} \right)^{1.5} \left(\frac{t}{t_i} \right)^{1.5m} \left(\frac{T}{10^4 \text{ K}} \right)^{0.68}, \end{aligned} \quad (7)$$

where

$$\langle \tau_{\text{eff}}^{0.5} \rangle = 0.67 [(\tau_{\text{CSM}_{\text{homogeneous}}} + \tau_{\text{CSM}_{\text{clumps}}})^{1.5} - \tau_{\text{CSM}_{\text{homogeneous}}}^{1.5}] \tau_{\text{CSM}_{\text{clumps}}}^{-1} \quad (8)$$

This estimate depends on the parameterized model as well as on assumptions of temperature ($T_{\text{CSM}} = 2 \times 10^4 \text{ K}$), pre-SN wind speed ($w_{\text{wind}} = 10 \text{ km s}^{-1}$), shock velocity ($v_i = 10^4 \text{ km s}^{-1}$), and t_i which accounts for the time between the appearance of optical lines from which the expansion velocity is determined, and the explosion date of the SN, typically assumed to be 45 days (Weiler *et al.* 1986). Equation 7 is derived from Case 2⁶ presented by Weiler *et al.* (2001; 2002). Since the SN was detected at such a late time, we can only approximately estimate the shock velocity and temperature at early times, making the mass loss rate estimates for SN 2001gd rather uncertain. However, the estimated mass loss rate of $3 \times 10^{-5} M_{\odot} \text{ yr}^{-1}$ is close to the values found for both SNe 1993J and 1985L. As one might expect since they are both type IIb SNe, there are many similarities between SNe 2001gd and 1993J, especially in that they require the presence of both uniform and clumpy (or filamentary) media to account for the radio light curve behavior while the mass

⁶Case 1 is a uniform, external absorbing medium; Case 2 is a clumpy or filamentary medium with a statistically large number of clumps; and Case 3 is similar to Case 2, but with a statistically small number of clumps.

loss rate for SN 2001gd appears to be in the lower range of known mass loss rates for type II SNe (1×10^{-5} — $1 \times 10^{-4} M_{\odot} \text{ yr}^{-1}$), it clearly exceeds the values for typical Ib/Ic SNe (7×10^{-7} — $1 \times 10^{-5} M_{\odot} \text{ yr}^{-1}$; Weiler *et al.* 2002).

4. Conclusions

We present new observations of SN 2001gd at multiple radio wavelengths for the first year after explosion. Modeling implies that the evolution of the radio emission from SN 2001gd behaves in a systematic fashion and is consistent with the non-thermal emitting, thermal absorbing model previously used successfully for other RSNe. The CSM required by the model fits is best described as a “filamentary” or “clumpy” medium embedded in a uniform component and appears similar to that suggested for the type IIb SN 1993J (Van Dyk *et al.* 1994). In general, SN 2001gd is very similar to SN 1993J suggesting that type IIb SNe may be relatively uniform in their properties. By comparison, the type II-L SN 1985L in the same galaxy as SN 2001gd appears significantly different in its radio properties, illustrating the general diversity of type II RSNe.

We are indebted to Dr. Barry Clark at the VLA for scheduling our numerous observations and to observers who contributed their telescope time. KWW wishes to thank the Office of Naval Research (ONR) for the 6.1 funding supporting this research. CJS is a National Research Council Postdoctoral Fellow. We would like to thank A. Ray and P. Chandra for sharing their 21 cm GMRT observation. Additional information and data on radio supernovae can be found on <http://rsd-www.nrl.navy.mil/7214/weiler/> and linked pages. We also would like to thank our anonymous referee for his/her useful comments.

REFERENCES

- Aksenov, E. P., Terebizh, Y. V., Metlova, N., Leibundgut, B., & Albert, T. 1985, IAU Circ., 4077
- Chandra, P., Ray, A., & Bhatnagar, S. 2002, IAU Circ.7982
- Chevalier, R. A. 1982a, ApJ, 259, 302
- Chevalier, R. A. 1982b, ApJ, 259, L85
- Chevalier, R. A. 1998, ApJ, 499, 810

- Filippenko, A. V., & Sargent, W. L. W. 1985, IAU Circ. 4080
- Kriss, G. 1985, IAU Circ. 4080
- Kowal, C. T., Sargent, W. L. W., & Zwicky, F. 1970, PASP, 82, 736
- Matheson, T., Jha, S., Challis, P., Kirshner, R., & Berlind, P. 2001, IAU Circ. 7765
- Montes, M. J., Weiler, K. W., & Panagia, N. 1997, ApJ, 488, 792
- Natta, A., & Panagia, N. 1984, ApJ, 287, 228
- Nakano, S., Itagaki, K., Kushida, Y., Kushida, R., & Dimai, A. 2001, IAU Circ. 7761
- Panagia, N., Sramek, R. A., & Weiler, K. W. 1986, ApJ, 300, L55
- Readhead, A. C. S. 1994, ApJ, 426, 51
- Schlegel, E. M., Ryder, S., Staveley-Smith, L., Petre, R., Colbert, E., Dopita, M., & Campbell-Wilson, D. 1999, AJ, 118, 2689
- Sramek, R. A., Panagia, N., & Weiler, K. W. 1984, ApJ, 285, L59
- Stockdale, C. J., Perez-Torres, M. A., Marcaide, J. M., Sramek, R. A., Weiler, K. W., Van Dyk, S. D., Panagia, N., Lundqvist, P., Pooley, D., Immler, S., & Lewin, W. 2002, IAU Circ. 7830
- Tully, R. B. 1988, Nearby Galaxies Catalog (Cambridge and New York: Cambridge University Press)
- Van Dyk, S. D., Weiler, K. W., Sramek, R. A., Rupen, M., & Panagia, N. 1994, ApJ, 432, L115
- Van Dyk, S. D., Montes, M. J., Weiler, K. W., Sramek, R. A., & Panagia, N. 1998, AJ, 115, 1103
- Weiler, K. W., Sramek, R. A., Panagia, N., van der Hulst, J. M., & Salvati, M. 1986, ApJ, 301, 790
- Weiler, K. W., Panagia, N., & Sramek, R. A. 1990, AJ, 364, 611
- Weiler, K. W., Panagia, N., & Montes, M. J. 2001, ApJ, 562, 670
- Weiler, K. W., Panagia, N., Montes, M. J., & Sramek, R. A. 2002, *Ann. Rev. Astron. Astrophys.*, 40, 387

Table 1. Calibrator Measurements for J1310+323

Obs. Date	Ref. Epoch (days)	Tel. Config.	S(21cm) (Jy)	S(6cm) (Jy)	S(3.6cm) (Jy)	S(2cm) (Jy)	S(1.2cm) (Jy)
2001-Nov-24	$\equiv 0.00$						
2002-Feb-08	75.72	VLA-A	1.703	1.659	1.765		2.145
2002-Mar-02	97.60	VLA-A	1.703 ^a	1.687	1.745	2.027	2.164
2002-Mar-14	109.66	VLA-A	1.700				
2002-Mar-23	118.42	VLA-A	1.681		1.798 ^a	2.077	2.315
2002-Apr-06	132.38	VLA-A	1.690	1.700 ^a	1.833 ^a	2.073	
2002-Jun-12	199.30	VLA-B	1.668	1.726	2.005		
2002-Jul-16	234.17	VLA-B	1.722	1.773	2.045		
2002-Oct-28	337.15	VLA-C			2.130	2.551	2.728

^aInterpolated from other calibrator measurements.

Table 2. Flux Density Measurements for SN 2001gd

Obs. Date	Ref. Epoch (days)	Tel. Config.	S(21cm) $\pm \sigma_f$ (mJy)	S(6cm) $\pm \sigma_f$ (mJy)	S(3.6cm) $\pm \sigma_f$ (mJy)	S(2cm) $\pm \sigma_f$ (mJy)	S(1.2cm) $\pm \sigma_f$ (mJy)
2001-Nov-24 ^a	$\equiv 0.00$						
2002-Feb-08	75.72	VLA-A	0.848 ± 0.129	5.050 ± 0.257	4.090 ± 0.208		1.370 ± 0.176
2002-Mar-02	97.60	VLA-A	1.414 ± 0.177	5.414 ± 0.284	3.732 ± 0.195	2.183 ± 0.242	1.335 ± 0.182
2002-Mar-14	109.66	VLA-A	1.840 ± 0.352		3.660 ± 0.208	2.330 ± 0.322	1.000 ± 0.187
2002-Mar-23	118.42	VLA-A	1.830 ± 0.339		3.780 ± 0.212	2.660 ± 0.304^b	
2002-Apr-06	132.38	VLA-A	1.990 ± 0.344	5.330 ± 0.277	3.100 ± 0.168		
2002-Jun-12	199.30	VLA-B	3.230 ± 0.365	4.842 ± 0.249			
2002-Jul-16	234.17	VLA-B	2.733 ± 0.314	4.260 ± 0.234	2.390 ± 0.137		
2002-Sep-22	301.40	GMRT	3.550 ± 0.220^c				
2002-Oct-28	337.15	VLA-C			2.071 ± 0.136	0.893 ± 0.178	0.809 ± 0.155

^aNakano *et al.* (2001); Matheson *et al.* (2001); SN 2001gd is believed to have exploded roughly 83 days prior to its initial optical detection. See Section 1 for further explanation.

^bThis value was not included in the modeling.

^cChandra, Ray, & Bhatnagar (2002)

Table 3. Fit of SN 2001gd & Comparison with other Type II Supernovae.

Parameter ^a	SN 2001gd	SN 1993J ^b	SN 1985L ^c
K_1	$(1.49^{+1.88}_{-1.09}) \times 10^3$	4.14×10^3	4.8×10^3
α	$-1.38^{+0.13}_{-0.12}$	-0.99	> -0.58
β	$-0.96^{+0.24}_{-0.13}$	-0.64	< -0.71
K_2	$(3.25^{+4.55}_{-1.31}) \times 10^6$	1.35×10^3	$> 7.00 \times 10^6$
δ	$\equiv -3.00$	-1.99	$\equiv -3.00$
K_3	$(1.05^{+1.31}_{-0.95}) \times 10^3$	2.54×10^4	0.00
δ'	$-1.27^{+0.40}_{-0.15}$	-2.02	0.00
Time to $L_{6 \text{ cm peak}}$ (days)	173^{+10}_{-15}	168	$\equiv 30$
$L_{6 \text{ cm peak}}$ (erg s ⁻¹ Hz ⁻¹)	$(2.91^{+0.13}_{-0.04}) \times 10^{27}$	1.45×10^{27}	$\sim 3.1 \times 10^{26}$
$\dot{M}$ (M _⊙ yr ⁻¹)	3.0×10^{-5}	2.1×10^{-5}	$> 6.3 \times 10^{-5}$

^aAssumes a constant mass-loss rate, constant velocity, wind-established CSM, i.e., $\rho \propto r^{-2}$.

^bVan Dyk *et al.* (1994)

^cVan Dyk *et al.* (1998)

Fig. 1.— Plot of the flux density of the VLA calibrator source J1310+323 at 1.3 cm (*circles*), 2 cm (*crosses*), 3.6 cm (*squares*), 6 cm (*triangles*), and 21 cm (*diamonds*). Note that the values for 21 cm are all reduced by 0.5 Jy to avoid confusion on the plot.

Fig. 2.— The radio measurements and parameterized model for SN 2001gd at 1.3 cm (*circles; solid line*), 2 cm (*crosses; dashed line*), 3.6 cm (*squares; dash-dot line*), 6 cm (*triangles; dotted line*), and 21 cm (*diamonds; dash-triple dotted line*). The lines represent the best fit model as described in the text with the parameters listed in Table 2. SN 2001gd is believed to have exploded 83 days prior to its initial optical discovery, see Section 1 for further explanation (Nakano *et al.* 2001; Matheson *et al.* 2001).



